

**Ex 1.**

- (1) Write down all Brauer tree with 5 edges and trivial exceptional multiplicity.
- (2) Write down the Cartan matrix of each of the associated Brauer tree algebra.

**Ex 2.** Let  $(T, v_0, m_0 = 1)$  be the multiplicity-free Brauer star with 3 edges.

- (1) Write down quiver  $Q_{T, v_0, m_0}$  and the admissible ideal  $I_{T, v_0, m_0}$ .
- (2) Write down all 9 strings of  $(Q_{T, v_0, m_0}, I_{T, v_0, m_0})$ ; you can write the associated module diagram instead if you prefer.

**Ex 3.** Repeat the previous question for the multiplicity-free Brauer line with 3 edges.

**Ex 4.** Consider an arrow  $(y|x)_v \in Q_T$  for a Brauer tree  $(T, v_0, m_0)$ . Let  $B$  be the associated Brauer tree algebra. Define a module  $H_{x,v} := B(y|x)_v$ . Note that when  $x$  is a leaf attached to valency 1 vertex  $v$ , then  $H_{x,v}$  is the simple module  $S_x$ .

- (1) Show that the module  $H_{x,v}$  is uniserial and find the corresponding string.
- (2) There is a surjective homomorphism  $P_x \rightarrow H_{x,v}$  with kernel  $H_{y,u}$ . What is  $y$  and  $u$ ?

**Ex 5.** Recall the construction of trivial extension algebra  $\Lambda \ltimes D\Lambda$  from Lecture 8. Let  $\Lambda$  be the lower triangular matrix ring

$$\begin{pmatrix} K & 0 \\ K & K \end{pmatrix} \cong K(1 \xrightarrow{\alpha} 2).$$

Note that  $D\Lambda \cong \begin{pmatrix} K & K \\ 0 & K \end{pmatrix}$  as (left or right or bi-) module. Show that the induced trivial extension algebra is isomorphic to the multiplicity-free Brauer tree algebra with 2 edges

$$B(T, v_0, m_0 = 1) = K(1 \begin{matrix} \xrightarrow{\alpha} \\ \xleftarrow{\alpha^*} \end{matrix} 2) / (\alpha\alpha^*\alpha, \alpha^*\alpha\alpha^*).$$

Deadline: 30th January, 2023