

You may assume all algebras are finite-dimensional over a field \mathbb{k} . You may attempt the exercises with the additional assumption of \mathbb{k} being algebraically closed.

Throughout, unadorned tensor product over assumed to be taken over a field, i.e. $\otimes = \otimes_{\mathbb{k}}$. For a module X over some algebra, denote by $\mathbf{add}(X)$ the full subcategory of the module category consisting of finite direct sums of direct summands of X (up to isomorphism).

Ex 1. Let e be an idempotent of a finite-dimensional algebra A . Consider the functors

$$\begin{array}{ccc} & \xleftarrow{\mathbf{l} := - \otimes_{eAe} eA} & \\ \mathbf{mod} A & \xrightleftharpoons[\mathbf{r} := \mathrm{Hom}_{eAe}(Ae, -)]{\mathbf{j} := - \otimes_A Ae} & \mathbf{mod} eAe \\ & \xrightarrow{\mathbf{r} := \mathrm{Hom}_{eAe}(Ae, -)} & \end{array}$$

- (1) Show that there are natural isomorphisms $\mathbf{j}\mathbf{r} \cong \mathrm{Id}_{\mathbf{mod} eAe} \cong \mathbf{j}\mathbf{l}$.
- (2) Show that $\mathbf{l}(P) \in \mathbf{add}(eA)$ for all projective eAe -module P .
Hint: Consider first the case when $P = fAe$ is an indecomposable projective eAe -module, where f is some primitive idempotent of A .
*** If you only present this as a consequence of property of adjointness, no mark will be awarded.*
- (3) For $N \in \mathbf{mod} eAe$, show that $M := \mathbf{l}(N) = N \otimes_{eAe} eA$ satisfies the following condition:

$$\text{There is an exact sequence } P_1 \xrightarrow{f} P_0 \xrightarrow{\pi} M \rightarrow 0 \text{ with } P_1, P_0 \in \mathbf{add}(eA). \quad (\dagger)$$
- (4) Let $M \in \mathbf{mod} A$. Show that M satisfies (\dagger) implies that $\mathbf{l}\mathbf{j}(M) = Me \otimes_{eAe} eA \cong M$.
Hint: Use (2) and find an appropriate commutative diagram.
- (5) Show that \mathbf{l}, \mathbf{j} defines an equivalence of categories $\mathbf{pres}(eA) \simeq \mathbf{mod} eAe$, where $\mathbf{pres}(eA)$ is the full subcategory of $\mathbf{mod} A$ consisting of modules M satisfying (\dagger) .

Ex 2.

- (1) Show that $\mathrm{Hom}_A(M, N) \cong D(M \otimes_A DN)$ as vector spaces.
- (2) Let $P_{\bullet} = (P_i, d_i : P_i \rightarrow P_{i-1})_{i \geq 0}$ be a projective resolution of an A -module M , and define

$$\mathrm{Tor}_1^A(M, N) := H_1(P_{\bullet} \otimes_A N) = \frac{\mathrm{Ker}(d_1 \otimes_A N)}{\mathrm{Im}(d_2 \otimes_A N)}$$

the first homology group of the complex $P_{\bullet} \otimes_A N$. Show that $\mathrm{Ext}_A^1(M, N) \cong D \mathrm{Tor}_1^A(M, DN)$ as \mathbb{k} -vector spaces.

- (3) Show that $D \mathrm{Hom}_A(M, A) \cong M \otimes_A DA$ as right A -modules.
- (4) Let ${}_A X_B$ be an A - B -bimodule. If M is a C - A -bimodule and N is a C - B -bimodule. Show that $\mathrm{Hom}_{C^{\mathrm{op}} \otimes B}(M \otimes_A X, N) \cong \mathrm{Hom}_{C^{\mathrm{op}} \otimes A}(M, \mathrm{Hom}_B(X, N))$ as vector spaces.

- (5) Let $B := A^{\text{op}} \otimes A$. Show that $\text{Hom}_B(A, B) \cong \text{Hom}_A(DA, A)$ as A - A -bimodules.
Hint: $B \cong (DDA) \otimes A \cong \text{Hom}_{\mathbb{k}}(DA, A)$ as B -modules.
- (6) In the setup of (5), show that $\text{Ext}_B^1(A, B) \cong \text{Ext}_A^1(DA, A)$.

Ex 3. Consider the quiver algebra $A = \mathbb{k}Q/I$ given by

$$Q : 1 \begin{array}{c} \xrightarrow{\alpha_1} \\ \xleftarrow{\beta_1} \end{array} 2 \begin{array}{c} \xrightarrow{\alpha_2} \\ \xleftarrow{\beta_2} \end{array} 3 \begin{array}{c} \xrightarrow{\alpha_3} \\ \xleftarrow{\beta_3} \end{array} 4, \quad I = (\beta_3\alpha_3, \alpha_i\alpha_{i+1}, \beta_{i+1}\beta_i, \beta_i\alpha_i - \alpha_{i+1}\beta_{i+1} \mid i = 1, 2)$$

For $i \in \{1, 2, 3, 4\}$, let $\Delta(i) := P_i/\alpha_i A$ (with $\alpha_4 := 0$ as a convention).

- (1) Describe the Loewy filtration of each indecomposable projective A -module $P(i)$ with $1 \leq i \leq 4$. In particular, show that each of these has a simple socle, i.e. $\text{soc}P(i) \cong S(j_i)$ for some $1 \leq j_i \leq 4$.
- (2) Write down the minimal projective resolution of $\Delta(1)$ and determine its projective dimension.
- (3) Show that $\text{Ext}_A^k(\Delta(i), \Delta(j)) = 0$ whenever $i > j$ for any $k \geq 0$.
- (4) Show that $\text{Ext}_A^k(\Delta(i), \Delta(j)) = 0$ whenever $k > 3$ for any i, j .
- (5) Compute $\dim_{\mathbb{k}} \text{Ext}_A^k(\Delta(i), \Delta(j))$ for all possible i, j, k . Show your working.

Deadline: 19th December, 2025

Submission: In lecture or E-mail to (replace at by @) aaron.chan at math.nagoya-u.ac.jp